ST.JUDE
SPECIALTY PHARMACY
& MEDICAL SUPPLIES

IVIG PRESCRIPTION REFERRAL FORM

121 St. Nicholas Ave Brooklyn, NY 11237
TEL: 718-381-5116 FAX: 718-417-3621
Proudly serving over 30 years

Today's Date

☐ NEW PATIENT ☐ CURRENT PATIENT

Patient Name First Name Middle Name Last Name DOB Weight ☐ Male ☐ Female
Street Address Apt # City State Zip
Daytime Tel Evening Tel Cell Email
Ship to Patient at ☐ Home ☐ Work **OR** Patient will pick up at ☐ Physician Office ☐ Pharmacy Date Needed
DIAGNOSIS ☐ Polymiositis (710.4) ☐ CIDP & Immune Neuropathies w/ Paraproteinemia (273.1, 357.81) ☐ CVID (279.3) ☐ CIDP (357.81)
☐ Guillian-Barre Syndrome (357.0) ☐ Immune Neuropathy other than CIDP w/o Paraproteinemia (357.81) ☐ Vasculitis Neuropathy (357.1) ☐ Dermatomyositis (710.3)
☐ Lambert–Eaton Myasthenic Syndrome (358.1) ☐ Myathesnia Gravis (358.1) ☐ Diabetic Proximal Neuropathy (358.1) ☐ Other

Insured's Name Relation to Patient Eligible for Medicare ☐ Yes ☐ No If yes, Medicare#
Prescription Card ☐ Yes ☐ No If Yes, Carrier Tel Fax Policy/Group#
Bin# Pcn# RXID# RX Group#

Prescriber's Name Office Contact
Street Address Suite # City State Zip
Tel Fax Email
License# NPI# UPIN# DEA#

PRESCRIPTION

PLEASE ATTACH COPIES OF PATIENT'S INSURANCE CARDS

Does patient already have a line? ☐ Yes ☐ No If yes, type of line
IVIG to be infused via existing line: ☐ Yes ☐ No
First IVIG Infusion?
☐ Yes If yes, IgA level is more than 5 mg/dl: ☐ Yes ☐ No ☐ Not Available
Ig Quantification: IgA, IgG, IgM (prior to 1st IVIG infusion)
☐ No If no, brand/dose of IVIG: Last infusion Date:
Note: IVIG contains IgA and is contraindicated in/gA deficient patients with antibodies against IgA and history of hypersensitivity.

IVIG (IMMUNOGLOBIN) ORDER: *(IVIG brand will be chosen if not specified)*

IVIG DOSE: g/kg = g (round to nearest vial size) infuse intravenously (Range: 0.2-2 g/kg)
☐ Repeat dose **daily** x consecutive days total, repeat dose: monthly x months
☐ Repeat dose **weekly** x weeks total
☐ Repeat dose **monthly** x months total
☐ Other

SUGGESTED RATE OF INFUSION:
☐ 30 -150 ml/hr as tolerated (Increase rate gradually every 30 min by 20-30 ml/hr)
☐ Other

PRE-MEDICATIONS: TO BE ADMINISTERED 30 MIN PRIOR TO IVIG INFUSION (QTY:PER INFUSION):
☐ Diphenhydramine 25 - 50 mg PO Dispense:#2 (25 mg)
☐ Acetaminophen 650 mg PO Dispense:#2 (325 mg)
☐ Other QTY: QS

IN THE EVENT OF ANAPHYLAXIS:
- Stop Infusion and call MD & 911
- Diphenhydramine 25 - 50 mg IVP every 4 hours prn (Not to exceed 25 mg/min) QTY: 3 (50 mg)
- Epinephrine (1:1000) 0.4 mg SQ prn anaphylaxis, may repeat every 20 minutes x 2 QTY: 3 amp

Other

SUPPLIES FOR INFUSION (If Necessary)
☐ NaCl 0.9% / D5W for flush: flush Line/Port with (3 - 5 ml for PIV and 5-10 ml for Central Line/Port) per nursing agency protocol *(NaCl 0.9% / D5W will be used based on IVIG compatibility)*
☐ Heparin for flush (100 Units / ml) (if RN keeps PIVor if needed for Central Line), flush with 3-5 ml per nursing agency protocol
☐ Sterile water for reconstitution of powder to make the requested concentration (for Carimune NF) ☐ Other:

Prescriber's Signature (signature required. NO STAMPS) **Date**

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately. PLEASE NOTE: St. JUDE can only accept original prescription drug orders from patients, faxed prescriptions can be accepted only from the prescribing practitioners.